

*Disclaimer: This is a machine generated PDF of selected content from our products. This functionality is provided solely for your convenience and is in no way intended to replace original scanned PDF. Neither Cengage Learning nor its licensors make any representations or warranties with respect to the machine generated PDF. The PDF is automatically generated "AS IS" and "AS AVAILABLE" and are not retained in our systems. CENGAGE LEARNING AND ITS LICENSORS SPECIFICALLY DISCLAIM ANY AND ALL EXPRESS OR IMPLIED WARRANTIES, INCLUDING WITHOUT LIMITATION, ANY WARRANTIES FOR AVAILABILITY, ACCURACY, TIMELINESS, COMPLETENESS, NON-INFRINGEMENT, MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Your use of the machine generated PDF is subject to all use restrictions contained in The Cengage Learning Subscription and License Agreement and/or the Gale Health and Wellness Terms and Conditions and by using the machine generated PDF functionality you agree to forgo any and all claims against Cengage Learning or its licensors for your use of the machine generated PDF functionality and any output derived therefrom.*

## Three Stages to COVID-19 Brain Damage Identified by Top Neurologists in Journal of Alzheimer Disease Paper Baseline MRIs Encouraged for COVID Patients.

**Date:** June 28, 2020

**From:** Medical Letter on the CDC & FDA

**Publisher:** NewsRX LLC

**Document Type:** Article

**Length:** 794 words

Full Text:

2020 JUN 28 (NewsRx) -- By a News Reporter-Staff News Editor at Medical Letter on the CDC & FDA -- The Journal of Alzheimer's Disease has just published a paper with a comprehensive review of the COVID-19's effect on the nervous system which classifies brain damage caused by COVID-19 into three stages. One of the authors, nationally-recognized neurologist Dr. Majid Fotuhi, MD, PhD, who is the medical director of NeuroGrow Brain Fitness Center in Northern Virginia and an affiliate staff at Johns Hopkins Medicine, encourages the adoption of this three-stage classification, calls for more research on COVID's long-term effects on the brain, and stresses the need for patients to receive a brain MRI before leaving the hospital.

"We are learning that a significant number of hospitalized COVID-19 patients have various degrees of brain impairment. As a medical community, we need to monitor these patients over time as some of them may develop cognitive decline, attention deficit, brain fog, or Alzheimer's disease in the future. There is a lot we can do to promote brain healing in COVID-19 patients, but first we must understand the nature and severity of their neurological deficits. At the patient level, getting a baseline MRI before leaving the hospital is imperative so that we have a starting point to evaluate and treat them," explained Fotuhi.

In the just published paper, Dr. Fotuhi and his colleagues warn about neurological issues in patients who suffer from COVID-19, including stroke, seizures, confusion, dizziness, paralysis, and/or coma. Already, two dozen case reports are revealing the impact of COVID-19 on the brains of patients. In fact, one study from Wuhan, China, showed that 45% of patients with severe COVID-19 illness experience marked neurological deficits. Another study from France showed 84% of ICU patients with COVID-19 have positive abnormalities on their neurological examination, and that 15% of patients who leave the ICU have residual "dysexecutive function," which involves poor attention and difficulty with decision-making and controlling behavior.

The paper proposes the adoption of a three stage "NeuroCovid" classification scheme to provide a basis from which to build on future hypotheses and investigations regarding SARS-Cov2 and the nervous system. These stages include: NeuroCovid Stage I: The virus damage is limited to epithelial cells of nose and mouth and the main symptoms include transient loss of smell and taste. NeuroCovid Stage II: The virus triggers a flood of inflammation, called cytokine storm, which begins in the lungs and travels in the blood vessels throughout all body organs. This cytokine storm leads to the formation of blood clots which cause small or large strokes in the brain. NeuroCovid Stage III: An explosive level of cytokine storm damages the blood brain barrier, the protective insulation layer in blood vessels of the brain. As a result, blood content, inflammatory markers, and virus particles invade the brain and patients develop seizures, confusion, coma, or encephalopathy.

Fotuhi points out that many patients with COVID-19 may have no noticeable neurological symptoms at first; but in some cases, patients may present with neurological symptoms even before they have fever, cough, or shortness of breath. In addition to having an MRI while at the hospital, he stresses that patients will need to be monitored in a few months after their hospitalization.

"Our experience with previous forms of coronaviruses suggest that in the long-term patients may develop depression, insomnia, Parkinson's disease, memory loss, or accelerated aging in the brain," elaborated Fotuhi. "For those recovering from COVID-19, I recommend regular exercise, eating a heart healthy diet, reducing stress, and improving sleep; these are critical ways patients can rejuvenate their brain and minimize having poor outcomes in the future."

These interventions, along with targeted brain training and neurofeedback therapy, are the main features of Dr. Fotuhi's 12-week Brain Fitness Program. As published in the Journal of Prevention of Alzheimer's Disease (2016), 84% of elderly with cognitive impairment who complete this brain rehabilitation program gain improvements in their brain function and many of them experience growth in the parts of their brain for learning and memory. These findings were similar for patients who gained recovery from their

persistent post-concussion syndrome. The program will now be tailored for patients suffering from post-COVID neurological issues.

A Harvard- and Johns Hopkins-trained neurologist and neuroscientist, Dr. Fotuhi is widely regarded as an authority in the field of memory, Alzheimer's Disease, concussion treatment, ADHD, and increasing brain vitality at any age. View source version on businesswire.com: <https://www.businesswire.com/news/home/20200610005749/en/>

Keywords for this news article include: Viral, COVID-19, Dementia, Virology, Cytokines, Hospitals, SARS-CoV-2, RNA Viruses, Tauopathies, Alzheimer Disease, Health and Medicine, Risk and Prevention, Brain Diseases and Conditions, NeuroGrow Brain Fitness Center, Neurodegenerative Diseases and Conditions, Intercellular Signaling Peptides and Proteins.

Our reports deliver fact-based news of research and discoveries from around the world. Copyright 2020, NewsRx LLC

**Copyright:** COPYRIGHT 2020 NewsRX LLC

<http://www.newsrx.com/newsletters/Medical-Letter-on-the-CDC-and-FDA.html>

**Source Citation** (MLA 8th Edition)

"Three Stages to COVID-19 Brain Damage Identified by Top Neurologists in Journal of Alzheimer Disease Paper Baseline MRIs Encouraged for COVID Patients." *Medical Letter on the CDC & FDA*, 28 June 2020, p. 1644. *Gale Health and Wellness*, [link.gale.com/apps/doc/A627202169/HWRC?u=mliin\\_b\\_bpublic&sid=bookmark-HWRC&xid=37d8bd79](http://link.gale.com/apps/doc/A627202169/HWRC?u=mliin_b_bpublic&sid=bookmark-HWRC&xid=37d8bd79). Accessed 1 July 2021.

**Gale Document Number:** GALE|A627202169